

CEFACLOR GH

Cefaclor Modified Release Tablets

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about **CEFACLOR GH**. It does not contain all the available information.

It does not take the place of talking to your doctor and pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking **CEFACLOR GH** against the benefits it is expected to have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine. You may need to read it again.

CEFACLOR GH tablets contain cefaclor monohydrate as the active ingredient.

What CEFACLOR GH is Used For

CEFACLOR GH is an antibiotic used to treat infections in different parts of the body caused by bacteria.

CEFACLOR GH will not work against infections caused by viruses such as colds or the flu.

CEFACLOR GH belongs to a group of antibiotics called cephalosporins, which are closely related to penicillins.

CEFACLOR GH works by killing the bacteria causing your infection or by stopping its growth.

Your doctor may have prescribed **CEFACLOR GH** for another reason. Ask your doctor if you have any questions about why **CEFACLOR GH** has been prescribed for you.

This medicine is available only with a doctor's prescription.

There is no evidence that **CEFACLOR GH** is addictive.

Before you start using CEFACLOR GH tablets

When you must not take it

Do not take CEFACLOR GH tablets if:

- **You have an allergy to cefaclor, other cephalosporins or any of the ingredients listed at the end of this leaflet.**

Some of the symptoms of an allergic reaction may include skin rash, itching, difficulty breathing, and swelling of the face, lips or tongue.

- **You have had a serious allergic reaction to penicillins**

- The packaging is torn or shows signs of tampering

- The expiry date on the pack has passed.

If you take this medicine after the expiry date has passed, it may not work as well (or it may make you feel sick).

Do not give **CEFACLOR GH** to children. The safety and efficacy of this product in children has not been established.

If you are not sure whether you should start using **CEFACLOR GH**, talk to your doctor.

Before you take it

You must tell your doctor:

1. If you have any type of allergic reaction to cephalosporin medicines, penicillin medicines or any other antibiotic medicines.

You may have an increased chance of being allergic to **CEFACLOR GH** if you are allergic to cephalosporins or penicillins.

2. If you have any allergies to any other medicines or any other substances, such as foods, preservatives or dyes

3. If you have or have ever had any other health problems/medical conditions, including:

- kidney disease
- severe bowel conditions/disease
- liver disease

4. If you are pregnant or intend to become pregnant

Your doctor will discuss the risks and benefits of using **CEFACLOR GH** during pregnancy.

5. If you are breast-feeding or plan to breast-feed.

Your doctor will discuss the risks and benefits of using **CEFACLOR GH** when breast-feeding.

If you have not told your doctor about any of the above, tell them before you start taking CEFACLOR GH.

Taking other medicines

Tell your doctor if you are taking any other medicines, including medicines that you buy without a prescription from your pharmacy, supermarket or health food shop. Some medicines may interfere with **CEFACLOR GH**. These include:

- probenecid, a medicine used to treat gout or to prolong the action of certain antibiotics.
- magnesium- or aluminium-containing antacids, medicines used to treat stomach upsets or stomach ulcers.

These medicines may be affected by **CEFACLOR GH**, or may affect how well it works. You may need different amounts of your medicine, or you may need to take different medicines. Your doctor or

pharmacist has more information on medicines to be careful with or avoid while taking **CEFACLOR GH**.

- nausea
- vomiting
- heartburn
- diarrhoea

How CEFACLOR GH is taken

How much to take

Your doctor will decide what dose and how long you will receive **CEFACLOR GH**. This depends on your infection and other factors, such as your weight.

How to take it

Swallow the prescribed dose of **CEFACLOR GH** tablets whole with a full glass of water. Do not cut, crush or chew the tablets.

Keep taking this medicine for the full time of treatment, even if you begin to feel better after a few days.

If you do not complete the full course of treatment prescribed by your doctor, the infection may not clear completely or your symptoms may return.

If you forget to take it

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to. Otherwise, take it as soon as you remember and then go back to taking it as you would normally. If you are not sure whether to skip the dose, talk to your doctor or pharmacist.

Do not take a double dose to make up for the dose that you missed.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor or Poisons Information Centre (telephone 13 11 26) or go to casualty at your nearest hospital if you think that you or anyone else may have taken too much CEFACLOR GH. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention. Keep telephone numbers of these places/services handy.

If you take too many **CEFACLOR GH** tablets, you may have:

While you are using CEFACLOR GH tablets

Things you must do

If the symptoms of your infection do not improve within a few days, or if they become worse, tell your doctor.

If you get severe diarrhoea tell your doctor, pharmacist or nurse immediately. Do this even if it occurs several weeks after **CEFACLOR GH** has been stopped.

Diarrhoea may mean that you have a serious condition affecting your bowel. You may need urgent medical care. Do not take any diarrhoea medicine without first checking with your doctor.

If you get a sore white mouth or tongue while taking or soon after stopping **CEFACLOR GH**, tell your doctor. Also tell your doctor if you get vaginal itching or discharge. This may mean you have a fungal infection called thrush. Sometimes the use of **CEFACLOR GH** allows fungi to grow and the above symptoms to occur. **CEFACLOR GH** does not work against fungi.

If you become pregnant while you are taking **CEFACLOR GH**, tell your doctor.

If you are about to start taking any new medicine, tell your doctor and pharmacist that you are taking **CEFACLOR GH**.

If you have to have any blood or urine tests, tell your doctor you are taking **CEFACLOR GH** tablets. **CEFACLOR GH** may affect the results of some blood and urine tests.

Tell all the doctors, dentists and pharmacists who are treating you that you are receiving **CEFACLOR GH**.

Things you must not do

Do not give CEFACLOR GH to anyone else, even if they have the same condition as you.

Do not use CEFACLOR GH to treat any other complaints unless your doctor tells you to.

Things to be careful of

Be careful driving or operating machinery until you know how **CEFACLOR GH** affects you.

CEFACLOR GH generally does not cause any problems with your ability to drive a car or operate machinery. However, as with many other medicines, **CEFACLOR GH** may cause dizziness or tiredness in some people.

Side Effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking CEFACLOR GH.

Like other medicines, **CEFACLOR GH** can cause some side effects. If they occur, most are likely to be minor and temporary. However some may be serious and need medical attention.

Ask your doctor or pharmacist to answer any questions you may have.

While you are taking it

Tell your doctor if you notice any of the following and they worry you:

- Oral thrush – white, furry, sore tongue and mouth
- Vaginal thrush – sore and itchy vagina and/or discharge
- Mild stomach upsets, such as feeling sick
- Diarrhoea

Tell your doctor immediately or go to casualty at your nearest hospital if you notice any of the following:

- fever
- skin rashes which may be itchy
- vomiting
- severe diarrhoea

These are all serious side effects of **CEFACLOR GH**. You may need urgent medical attention.

If any of the following happen, stop using CEFACLOR GH and tell your doctor immediately or go

to casualty at your nearest hospital:

- life threatening allergic reactions which may include skin rash, itching, difficulty breathing, swelling of the face, lips or tongue
- skin rash with joint pain and fever
- decrease in urine volume

These are all very serious side effects of **CEFACTOR GH**. If you have them, you may have had a serious allergic reaction to **CEFACTOR GH**. You may need urgent medical attention or hospitalisation.

After finishing it

Tell your doctor immediately if you notice any of the following side effects, particularly if they occur several weeks after stopping treatment with CEFACOR GH:

- severe abdominal cramps or stomach cramps
- watery and severe diarrhoea, which may also be bloody
- fever, in combination with one or both of the above

These are rare but serious side effects. You may have a serious condition affecting your bowel. **CEFACTOR GH** can cause bacteria which are normally present in the bowel and are normally harmless to multiply and thereby cause the above symptoms. You may need urgent medical attention. However this side effect is rare.

Do not take any diarrhoea medicine without first checking with your doctor.

Other side effects not listed above may also occur in some patients. Tell your doctor if you notice any other effects.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

After using CEFACOR GH tablets

Storage

At home, keep your **CEFACTOR GH** tablets in their original packs until it is time to use them. If you

take the tablets out of the pack, they may not keep well.

Keep it in a cool, dry place where the temperature stays below 25°C. Do not store it or any other medicine in the bathroom or near a sink.

Do not leave it in the car or on window sills.

Heat and dampness can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one and a half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking **CEFACTOR GH** or the tablets have passed their expiry date, ask your pharmacist what to do with any tablets that are left over.

Product description

What it looks like

CEFACTOR GH modified-release tablets 375 mg are capsule-shaped, biconvex, unscored, blue, film-coated tablets imprinted with "CEFACTOR CD 375 mg" in black ink. **CEFACTOR GH** tablets are available in a pack of 10 tablets.

Ingredients

Active ingredient:

Each tablet contains 375 mg cefaclor as cefaclor monohydrate.

Other ingredients:

Hypromellose, hydroxypropylcellulose, lactose, magnesium stearate, purified talc and colloidal anhydrous silica.

The film coating contains:

hypromellose, titanium dioxide, Macrogol 400 and indigo carmine CI73015.

The printing ink contains: purified water, iron oxide black, propylene glycol, hypromellose and isopropyl alcohol.

CEFACTOR GH does not contain gluten, sucrose, tartrazine or any other azo dyes.

Australian Registration No.

AUST R 182048

Sponsor

Ranbaxy Australia Pty Ltd
Suite 4.02, Level 4, Building D
12-24, Talavera Road
North Ryde, NSW 2113
Australia

CEFACTOR GH is distributed in Australia by:

Generic Health Pty Ltd
Suite 1, Level 1
1175 Toorak Road
Camberwell VIC 3124
Australia

Date of leaflet preparation: Jun 2011