



CONSUMER MEDICINE INFORMATION

Levetiracetam-GH Tablets 250 mg, 500 mg and 1000 mg

Levetiracetam

WHAT IS IN THIS LEAFLET

This leaflet answers some common questions about Levetiracetam-GH Tablets. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Levetiracetam-GH Tablets against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

WHAT LEVETIRACETAM-GH TABLETS IS USED FOR

Levetiracetam-GH Tablets is used to control epilepsy.

Epilepsy is a condition where you have repeated seizures. There are many different types of seizures, ranging from mild to severe.

This medicine belongs to a group of medicines called antiepileptics. These medicines are thought to work by controlling brain chemicals which send signals to nerves so that seizures do not happen.

Levetiracetam-GH Tablets may be used alone, or in combination with other medicines, to treat your condition.

Your doctor may prescribe Levetiracetam-GH Tablets in addition to your current therapy.

Ask your doctor if you have any questions about why Levetiracetam-GH Tablets has been prescribed for you.

There is no evidence that Levetiracetam-GH Tablets is addictive.

This medicine is available only with a doctor's prescription.

The safety and effectiveness of Levetiracetam-GH Tablets has not been established in patients less than 4 years of age.

BEFORE YOU TAKE LEVETIRACETAM-GH TABLETS

When you must not take it

Do not take Levetiracetam-GH Tablets if you have an allergy to:

- any of the ingredients listed at the end of this leaflet

Symptoms of an allergic reaction to Levetiracetam-GH Tablets may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin

Do not take this medicine after the expiry date (EXP) printed on the pack.

Do not take this medicine if the packaging is torn or shows signs of tampering or if the tablets do not look quite right. If this medicine has expired or is damaged, return it to your pharmacist for disposal.

If you are unsure whether you should start taking this

medicine, talk to your doctor or pharmacist.

Before you start to take it

Tell your doctor or pharmacist if you have allergies to:

- any other medicines, especially barbiturates (such as phenobarbitone) or any other antiepileptic medicines (such as carbamazepine, lamotrigine or valproate)
- any other substances, such as foods, preservatives or dyes.

Tell your doctor if you have or have had any medical conditions, especially the following:

- kidney problems
- liver problems

Tell your doctor if you are pregnant or intend to become pregnant.

Levetiracetam-GH Tablets may affect your developing baby if you take it during pregnancy. However, it is very important to control your seizures while you are pregnant. Your doctor will outline and weigh up all the risks and benefits of taking Levetiracetam-GH during pregnancy to help decide whether or not you should take it.

Tell your doctor if you are breast-feeding or plan to breast-feed.

The active ingredient in Levetiracetam-GH passes into breast milk and there is a possibility that your baby may be affected. Your doctor will discuss the risks and benefits of using Levetiracetam-GH if you are breastfeeding.

If you have not told your doctor or pharmacist about any of the above, tell them before you start taking Levetiracetam-GH Tablets.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.

Levetiracetam-GH Tablets does not interact with the oral contraceptive pill. However, you may be given Levetiracetam-GH Tablets together with other antiepileptic medicines that do interact and may affect the effectiveness of your contraceptive.

Your doctor may advise you to use an additional method of contraception if you take Levetiracetam-GH Tablets with other antiepileptic medicines.

**HOW TO TAKE
LEVETIRACETAM-GH
TABLETS**

How much to take

Follow all directions given to you by your doctor carefully. They may differ from the information contained in this leaflet.

Your doctor will tell you how much Levetiracetam-GH you will need to take each day. This may depend on your age, your condition and whether or not you are taking any other medicines.

For patients 12 years of age and older, the dosage is

generally between 1000mg and 3000mg each day. Taken in two doses.

For children 4 to 11 years of age the doctor will calculate the dosage based on the child's weight and tell you how much to give. The medicine is to be given twice daily.

Your doctor may start you on a low dose of Levetiracetam-GH Tablets first. Your doctor will slowly increase the amount of medicine until you are taking enough to control your epilepsy and you are not having seizures.

If you do not understand the instructions on the pack, ask your doctor or pharmacist for help.

How to take it

Swallow the tablets whole with a glass of water.

When to take it

Take Levetiracetam-GH Tablets twice a day, once in the morning and once at night. Take it, at about the same time each day.

Taking your medicine at the same time each day will have the best effect. It will also help you remember when to take it.

It does not matter if you take Levetiracetam-GH Tablets before or after food.

If you forget to take it

Contact your doctor if you have missed one or more doses.

Do not take a double dose to make up for the dose that you missed. This may

increase the chance of you getting an unwanted side effect.

If you are not sure what to do, ask your doctor or pharmacist.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

How long to take it

Most antiepileptic medicines take time to work, so do not be discouraged if you do not feel better straight away.

Continue taking your medicine for as long as your doctor tells you to. This medicine helps control your condition, but does not cure it. Therefore you must take your medicine every day, even if you feel well.

Do not stop taking Levetiracetam-GH Tablets, or change the dosage, without checking with your doctor. Do not let yourself run out of medicine over the weekend or on holidays. Stopping Levetiracetam-GH Tablets suddenly may cause unwanted effects or make your condition worse. Your doctor will slowly reduce your dose before you can stop taking it completely.

If you take too much (overdose)

Immediately telephone your doctor or the Poisons Information Centre (telephone 13 11 26) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have

taken too much Levetiracetam-GH Tablets. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Symptoms of An overdose may include feeling drowsy.

WHILE YOU ARE TAKING LEVETIRACETAM-GH TABLETS

Things you must do

Tell your doctor immediately if you notice an increase in seizures.

Tell your doctor immediately if you have symptoms of depression or thoughts of harming yourself.

Tell any other doctors, dentists, and pharmacists who are treating you that you are taking this medicine.

If you are about to be started on any new medicine, tell your doctor, dentist or pharmacist that you are taking Levetiracetam-GH Tablets.

Before you have any surgery or emergency treatment, tell your doctor or dentist that you are taking Levetiracetam-GH Tablets.

Tell your doctor if you feel Levetiracetam-GH Tablets is not helping your condition. Your doctor may need to change your medicine.

Tell your doctor if, for any reason, you have not taken this medicine exactly as prescribed. Otherwise, your

doctor may change your treatment unnecessarily.

If you become pregnant while taking Levetiracetam-GH Tablets, tell your doctor.

Be sure to keep all of your doctor's appointments so that your progress can be checked. Your doctor will check your progress and may want to take some tests from time to time. This helps to prevent unwanted side effects.

Things you must not do

Do not give Levetiracetam-GH Tablets to anyone else, even if their symptoms seem similar to yours or they have the same condition as you.

Do not take Levetiracetam-GH Tablets to treat any other complaints unless your doctor tells you to.

Things to be careful of

Be careful driving or operating machinery until you know how Levetiracetam-GH Tablets affects you. Children should be careful doing things like riding bicycles or climbing trees.

As with other antiepileptic medicines, Levetiracetam-GH Tablets may cause drowsiness in some people.

This is more frequent at the beginning of treatment or after an increase in the dose.

If you are feeling dizzy or drowsy do not drive, operate machinery, or do anything else that could be dangerous. Children should not ride a bike,

climb trees or do anything else that could be dangerous if they are feeling dizzy or drowsy.

Be careful when drinking alcohol while taking this medicine.

Combining Levetiracetam-GH Tablets and alcohol can make you more drowsy.

Your doctor may suggest you avoid alcohol while you are being treated with Levetiracetam-GH Tablets.

SIDE EFFECTS

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Levetiracetam-GH Tablets.

This medicine helps most people with epilepsy, but it may have unwanted side effects in a few people. All medicines can have side effects.

Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

Do not be alarmed by the following list of side effects. You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

If you get any side effects, do not stop taking Levetiracetam-GH Tablets without first talking to your doctor or pharmacist.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- dizziness

- feeling weak
- common cold
- upset stomach
- diarrhoea
- feeling tired, drowsy or sleepy

The above list includes the more common side effects of your medicine. They are usually mild and short-lived.

Tell your doctor as soon as possible if you notice any of the following:

- mood changes such as depression, nervousness, aggression, anger, anxiety, confusion, hallucination, irritability.
- feelings of depression
- upper respiratory tract infections
- weight loss.

The above list includes serious side effects that may require medical attention.

If any of the following happen, tell your doctor immediately or go to Accident and Emergency.

If you experience more frequent or more severe seizures (fits), or thoughts of self harm tell your doctor immediately or go to Accident and Emergency at your nearest hospital.

- thought of harming yourself
- more frequent or more severe seizures
- shortness of breath
- wheezing or difficult breathing
- swelling of the face, lips, mouth, throat or tongue or other parts of the body
- rash, itching or hives on the skin.

The above list includes very serious side effects. You may

need urgent medical attention or hospitalization.

Tell your doctor if you notice anything else that is making you feel unwell. Other side effects not listed above may happen in some people.

AFTER USING LEVETIRACETAM-GH TABLETS

Storage

Keep your tablets in the pack until it is time to take them. If you take the tablets out of their pack they will not keep well.

Keep your tablets in a cool dry place where the temperature stays below 30 °C.

Do not store Levetiracetam-GH Tablets or any other medicine in the bathroom or near a sink.

Do not leave it on a window sill or in the car on hot days. Heat and dampness can destroy some medicines. **Keep it where children cannot reach it.** A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

PRODUCT DESCRIPTION

What it looks like

Levetiracetam-GH Tablets are available in three strengths: 250mg -Blue, oblong -shaped, biconvex film-coated tablets, debossed with '250' on one side and score line on the other side.

500mg - yellow, oblong-shaped, biconvex film-coated tablets, debossed with '500' on one side and score line on the other side.

1000mg - White to off white, Oblong -shaped, biconvex film-coated tablets, debossed with '1000' on one side and score line on the other side.

Levetiracetam 250mg, 500mg and 1000mg tablets are packed in blister packs, pack sizes of 60 Tablets.

.Ingredients

Active ingredient

Each Levetiracetam-GH Tablet contains 250 mg, 500mg and 1000 mg of Levetiracetam as the active ingredient.

The inactive ingredients in Levetiracetam-GH include:

Starch - maize
Silica-colloidal anhydrous
Croscarmellose sodium
Povidone
Cellulose - Microcrystalline
Talc-purified
Magnesium stearate

Tablets are film-coated & the coating for each tablet strength contains polyvinyl alcohol, macrogol 6000 and macrogol 3350, talc - purified and titanium dioxide (E 171).

The following strengths also contain:

250mg - Indigo carmine
CI73015
500mg - Iron oxide yellow
CI77492

Levetiracetam-GH Tablets do not contain lactose, sucrose, gluten, tartrazine or any other azo dyes.

Levetiracetam-GH 250 mg -
AUST R 161294

Levetiracetam-GH 500 mg -
AUST R 161295

Levetiracetam-GH 1000 mg -
AUST R 161296

Supplier :

Levetiracetam-GH Tablets

is distributed in Australia
by:

Generic Health Pty Ltd.
Level 1, 1102 Toorak Road
Camberwell VIC 3124

Date of preparation

February 2014