

Pharmacy Action Pharmisil

Terbinafine Hydrochloride 1% cream

Pharmacy Action Pharmisil

This leaflet provides some useful information on your medicine. Please read it carefully before you start using this medicine. This leaflet does not contain all the information about your medicine.

If you have any further questions or are not sure about anything, ask your pharmacist.

What is Pharmacy Action Pharmisil

Pharmacy Action Pharmisil contains the active ingredient terbinafine hydrochloride. It also contains sorbitan monostearate, cetyl palmitate, cetyl alcohol, stearyl alcohol, polysorbate 60, isopropyl myristate, sodium hydroxide, purified water and benzyl alcohol.

What Pharmacy Action Pharmisil is used for

Pharmacy Action Pharmisil is used to treat tinea of the feet, body and groin region.

Tinea, commonly known as ringworm, is an infection of the skin caused by a type of fungi called dermatophytes. Tinea between the toes is commonly known as athlete's foot. Pharmacy Action Pharmisil is also used to treat candida (yeast) infection of the skin. Yeasts are another type of fungus.

Before using Pharmacy Action Pharmisil

Before using Pharmacy Action Pharmisil it is important to read the following:

What if I'm allergic to Pharmacy Action Pharmisil?

Pharmacy Action Pharmisil should not be used by people who are allergic to the drug terbinafine or to any other ingredient in the product.

Can Pharmacy Action Pharmisil be used on children?

Pharmacy Action Pharmisil may be used on children aged 12 years or older. Use on younger children is not recommended.

Should Pharmacy Action Pharmisil be used if I am pregnant or intend to become pregnant?

Since there is little experience with the safety of using Pharmacy Action Pharmisil in pregnancy, the cream should not be used by pregnant women or those who are trying to become pregnant.

Should Pharmacy Action Pharmisil be used if I'm breastfeeding?

You should not use Pharmacy Action Pharmisil if you are breastfeeding.

How to use Pharmacy Action Pharmisil

You should apply Pharmacy Action Pharmisil only to skin outside the body. Wash your hands after use. Avoid applying the cream near your eyes or touching your eyes while you still have the cream on your hands. In case of accidental contact with the eyes, rinse the eyes thoroughly with running water and if any symptoms persist see your doctor.

Apply Pharmacy Action Pharmisil once daily.

1. Clean and thoroughly dry the affected areas on your skin
2. Apply a sufficient amount of the cream on the affected skin and surrounding area. Use only a thin layer of cream and rub in gently.
3. If you need to apply Pharmacy Action Pharmisil between your toes, fingers or buttocks, under the breast or armpit, or in the groin region, cover the area with gauze after the cream has been applied. This is particularly useful at bedtime.

How long to use this medicine

How long you need to apply Pharmacy Action Pharmisil for depends on the type of infection you have:

Tinea (athlete's foot, ringworm, Jock itch)	7 days
Tinea infection on the sole of the foot	2-4 weeks
Candida (yeast) infections	1-2 weeks

While the signs of the infection may disappear after only using Pharmacy Action Pharmisil for a few days, it is important to continue use for the recommended period. The reason is that if you use it for only a few days, you may not remove all of the fungi from the affected areas.

And as a result the infection may return soon after you stop using the product.

If there is no improvement in your condition after using Pharmacy Action Pharmisil for the recommended time, see your doctor or pharmacist.

Side effects

Some people may experience redness, itchiness or a stinging sensation in the areas where PHARMACY ACTION Pharmisil has been applied.

Although these symptoms are normally harmless, you should nevertheless tell your pharmacist if they occur, as you may be experiencing an allergy to Pharmacy Action Pharmisil. If your pharmacist confirms you are suffering an allergy you should stop using the product.

In case of accidental poisoning

If Pharmacy Action Pharmisil is accidentally swallowed, particularly by a child, contact your doctor or the Poison Information Centre (Telephone: Australia 13 11 26; New Zealand 0800 764 766).

Storage

- Store below 30°C
- Store out of reach of children
- Do not use after the expiry date shown on the pack.

Tamper Evident Packaging.

Pharmacy Action Pharmisil is enclosed in tamper-evident packaging to help protect you from product tampering. If you find that the tube seal has been broken or is missing, do not use the product but return it to your pharmacist.

Where to go for further information

If you have any further questions about Pharmacy Action Pharmisil contact your pharmacist.

Sponsor

Generic Health Pty Ltd
Level 1, 1102 Toorak Road
Camberwell, Vic, 3124