

This card contains important safety information that you need to be aware of before you are given Nepexto and during treatment with Nepexto.

If you do not understand this information, please ask your doctor to explain it to you.

**Show this card to any doctor involved in your treatment. Keep this card with you for 2 months after the last Nepexto dose, since side effects may occur after your last dose of Nepexto.**

**See the Nepexto Package Leaflet for more information.**

**It is important that you and your doctor record the brand name and batch number of your medication.**

**Brand name:**

**Batch number:**



**Sponsor:**

Maxx Pharma Pty Ltd  
Melbourne, Victoria 3000  
Australia

**Distributed by:**

Generic Health Pty Ltd.

**Nepexto<sup>®</sup>**  
**etanercept**

**Patient Alert Card**

BC00627/01

279853

**Infections**

Nepexto may increase your risk of getting infections, which could be serious.

You should not use Nepexto:

- if you have an infection.
- if you develop symptoms suggestive of infections, such as fever, chills, persistent cough, mouth ulcers or sore throat.

Seek medical attention immediately.

If you are not sure, ask your doctor.

You should be evaluated for tuberculosis (TB) or any other infections before starting Nepexto.

Ask your doctor to record the dates and the results of your last screening for TB below:

Test: \_\_\_\_\_  
Date: \_\_\_\_\_  
Result: \_\_\_\_\_

Test: \_\_\_\_\_  
Date: \_\_\_\_\_  
Result: \_\_\_\_\_

Please ask your physician to list your other medicines that may increase your risk of infection.

Other Medicines

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

**Congestive Heart Failure**

If you have congestive heart failure, and you think your symptoms are becoming worse (eg. shortness of breath, persistent cough or fatigue, difficulty exercising, swelling in the legs or feet), or if you experience new onset of such symptoms, seek medical attention immediately.

**Use in Children and Adolescents**

Nepexto is not indicated for use in children or adolescents weighing less than 62.5kg.

**Other Information  
(please complete)**

Patient's Name

\_\_\_\_\_  
\_\_\_\_\_

Doctor's Name

\_\_\_\_\_  
\_\_\_\_\_

Doctor's Phone

\_\_\_\_\_  
\_\_\_\_\_

If you or anyone you know experiences any side effects after using Nepexto, report them to [medical@maxxpharma.com.au](mailto:medical@maxxpharma.com.au) or 1800 202 981 and speak to your treating physician.